

Europäisches Patentamt
European Patent Office
Office européen des brevets



(11) **EP 1 123 979 A1**

(12)

EUROPEAN PATENT APPLICATION

published in accordance with Art. 158(3) EPC

(43) Date of publication:
16.08.2001 Bulletin 2001/33

(51) Int Cl.7: **C12N 15/54, C12N 9/12,
C12N 1/21, C12P 7/66**

(21) Application number: **00954944.5**

(86) International application number:
PCT/JP00/05659

(22) Date of filing: **24.08.2000**

(87) International publication number:
WO 01/14567 (01.03.2001 Gazette 2001/09)

(84) Designated Contracting States:
**AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU
MC NL PT**
Designated Extension States:
AL LT LV MK RO SI

(30) Priority: **24.08.1999 JP 23756199**

(71) Applicant: **KANEKA CORPORATION**
Osaka-shi, Osaka 530-8288 (JP)

(72) Inventors:
• **MATSUDA, Hideyuki**
Matsue-shi, Shimane 690-0815 (JP)

- **KAWAMUKAI, Makoto**
Matsue-shi, Shimane 690-0823 (JP)
- **YAJIMA, Kazuyoshi**
Akashi-shi, Hyogo 673-0005 (JP)
- **IKENAKA, Yasuhiro**
Kobe-shi, Hyogo 651-2242 (JP)
- **HASEGAWA, Junzo**
Akashi-shi, Hyogo 674-0057 (JP)
- **TAKAHASHI, Satomi**
Kobe-shi, Hyogo 655-0851 (JP)

(74) Representative: **Gillard, Marie-Louise et al**
Cabinet Beau de Loménie
158, rue de l'Université
75340 Paris Cédex 07 (FR)

(54) **PROCESS FOR PRODUCING COENZYME Q 10?**

(57) The present invention has for its object to isolate a gene coding for the enzyme synthesizing coenzyme Q₁₀ side chain synthase from a fungal strain of the genus *Saitoella* and exploit it to advantage for the efficient microbial production of coenzyme Q₁₀.

The present invention provides;

a DNA having the nucleotide sequence shown under SEQ ID NO:1;

a DNA having a nucleotide sequence derived from

the nucleotide sequence of SEQ ID NO:1 by deletion, addition, insertion and/or substitution of one or a plurality of nucleotides and coding for a protein having decaprenyl diphosphate synthase activity; a DNA which hybridizes with the DNA having the nucleotide sequence of SEQ ID NO:1 under stringent conditions and codes for a protein having decaprenyl diphosphate synthase activity.

EP 1 123 979 A1

Description

TECHNICAL FIELD

5 [0001] The present invention relates to a process for producing coenzyme Q₁₀ for pharmaceutical and other uses. More particularly, the invention relates to a process for producing coenzyme Q₁₀ which comprises isolating a gene coding for the coenzyme Q₁₀ side-chain synthase, which is a key enzyme involved in the biosynthesis of coenzyme Q₁₀, i.e. decaprenyl diphosphate synthase, from a fungal strain of the genus Saitoella and introducing it into a host microorganism to let it elaborate coenzyme Q₁₀.

BACKGROUND ART

[0002] The conventional technology for commercial production of coenzyme Q₁₀ comprises isolating the coenzyme from a tobacco or other plant and modifying the length of its side chain by a synthetic technique.

15 [0003] While it is known that coenzyme Q₁₀ is produced by a broad spectrum of organisms ranging from microorganisms, such as bacteria and yeasts, to higher animals and plants, the method comprising culturing a microorganism and extracting coenzyme Q₁₀ from the microorganism is regarded as one of the most effective production methods and has actually been exploited commercially. However, the prior art methods are invariably poor in productivity, providing only low yields and/or involving time-consuming procedures.

20 [0004] The pathways for biosynthesis of coenzyme Q₁₀ in organisms are partly different between the prokaryotes and the eukaryotes but invariably comprise a complicated cascade of reactions involving many kinds of enzymes. However, these pathways are basically comprised of three fundamental steps, namely the step of synthesizing decaprenyl diphosphate as the precursor of the prenyl side-chain of coenzyme Q₁₀, the step of synthesizing p-hydroxybenzoic acid as the basis of the quinone ring of coenzyme Q₁₀, and the step of coupling these two compounds together and effecting a serial substituent transformation to complete coenzyme Q₁₀. Of these reactions, the reaction determining the length of the side-chain of coenzyme Q₁₀ and acknowledged to be the rate-determining step of its biosynthesis, i.e. the reaction catalyzed by decaprenyl diphosphate synthase, is considered to be the most important reaction. Therefore, in order that coenzyme Q₁₀ may be produced with good efficiency, it seems worthwhile to isolate the key gene involved in said biosynthesis, namely the gene coding for decaprenyl diphosphate synthase, and utilize it for enhanced production of the enzyme. As sources of the gene, fungi capable of producing coenzyme Q₁₀ in comparatively large amounts can be regarded as useful candidates.

30 [0005] Heretofore, genes coding for decaprenyl diphosphate synthase have been isolated from several kinds of microorganisms, such as Schizosaccharomyces pombe (JP09-173076A) and Gluconobacter suboxydans (JP10-57072A), etc., but the inherent coenzyme Q₁₀ productivity of these microorganisms cannot be considered high enough and neither an efficient cultural protocol for these microorganisms nor an efficient isolation and purification procedure has been established as yet. Therefore, there has been a standing demand for isolation of a coenzyme Q₁₀-encoding gene from a microorganism capable of highly producing a coenzyme Q₁₀.

35 [0006] Devoted to providing a solution to the above-mentioned production problems, the present invention has for its object to isolate a gene coding for the enzyme synthesizing the coenzyme Q₁₀ side chain from a fungal strain of the genus Saitoella and exploit it to advantage for the efficient microbial production of coenzyme Q₁₀.

DISCLOSURE OF THE INVENTION

45 [0007] To accomplish the above object, in the present invention, the key gene involved in the biosynthesis of coenzyme Q₁₀, namely the gene coding for decaprenyl diphosphate synthase, was isolated from a fungal strain of the genus Saitoella in the first place. Then, this gene was introduced and allowed to be expressed in a host microorganism, such as Escherichia coli, to thereby enable the host to produce coenzyme Q₁₀ with efficiency.

[0008] The inventors of the present invention made intensive investigations for isolating such genes coding for decaprenyl diphosphate synthase from fungal strains of the genus Saitoella capable of producing comparatively large amounts of coenzyme Q₁₀ and have succeeded in isolating said genes.

50 [0009] The present invention, therefore, is concerned with a DNA of the following (a), (b) or (c):

(a) a DNA having the nucleotide sequence shown under SEQ ID NO:1;

55 (b) a DNA having a nucleotide sequence derived from the nucleotide sequence of SEQ ID NO:1 by deletion, addition, insertion and/or substitution of one or a plurality of nucleotides and coding for a protein having decaprenyl diphosphate synthase activity;

(c) a DNA which hybridizes with the DNA having the nucleotide sequence of SEQ ID NO:1 under stringent conditions and codes for a protein having decaprenyl diphosphate synthase activity.

[0010] The present invention is further concerned with a protein of the following (d) or (e).

(d) a protein having the amino acid sequence shown under SEQ ID NO:2;

(e) a protein having an amino acid sequence derived from the amino acid sequence of SEQ ID NO:2 by deletion, addition, insertion and/or substitution of one or a plurality of amino acids and having decaprenyl diphosphate synthase activity.

[0011] The invention is further concerned with a DNA coding for this protein.

[0012] The present invention is further concerned with an expression vector containing said DNA. For the expression vector of the invention, various vector systems heretofore known can be utilized and, therefore, may for example be pNTS₁ as constructed by cloning the DNA having the sequence of SEQ ID NO:1 into the vector pUCNT for expression.

[0013] The present invention is further concerned with a transformant as constructed by transforming a host microorganism with said DNA. As the host microorganism for the invention, *Escherichia coli* can be used with advantage.

[0014] The invention is further concerned with a process for producing coenzyme Q₁₀, which comprises culturing said transformant in a culture broth and harvesting the coenzyme Q₁₀ produced and accumulated in the resulting culture.

[0015] The host microorganism for this process is not particularly restricted but may be *Escherichia coli* to mention a preferred example. The coenzyme Q produced by *Escherichia coli* is coenzyme Q₈ but the invention enables this microorganism to produce coenzyme Q₁₀.

[0016] The inventors made intensive investigations on the isolation of the enzyme gene from a fungal strain which belongs to the genus *Saitoella* and is capable of producing comparatively large amounts of coenzyme Q₁₀ and succeeded in acquiring a fragment of the particular gene by the PCR technique.

[0017] The inventors compared the sequence of the known gene coding for decaprenyl diphosphate synthase with the genes coding for polyprenyl diphosphate synthases, namely long-chain prenyl synthases which are analogous to said known enzyme gene but differ from the same in chain length and, for the region of high homology and synthesized various PCR primers. Using these primers in various combinations, they studied PCR conditions. As a result, they found by analysis of the gene sequence that when a PCR, using DPS-1 (5'-AAGGATCCTNYTNCAYGAYGT-3') and DPS-1 1AS (5'-ARYTGNADRAAYTCNCC-3') [in the above sequences, R means A or G; Y means C or T, and N means G, A, T or C] as primers, is carried out according to the protocol of heat-treatment at 94 °C × 3 minutes, followed by 40 cycles of 94 °C, 1 minute → 43 °C, 2 minutes → 72 °C, 2 minutes, a ca. 220 bp fragment of the enzyme gene can be amplified from the chromosome gene of *Saitoella complicata* IFO 10748, a fungus belonging to the genus *Saitoella*.

[0018] Then, to acquire the full length of this enzyme gene, the chromosome gene of *Saitoella complicata* IFO 10748 is digested with the restriction enzyme EcoRI and inserted into a λ phage vector to construct a recombinant phage library. After the plaque is transferred to a nylon membrane, the plaque hybridization is carried out using the labeled PCR fragment, whereby a clone having the full-length decaprenyl diphosphate synthase gene can be obtained.

[0019] Sequencing of the decaprenyl diphosphate synthase gene occurring in the above clone reveals that the gene has the nucleotide sequence shown under SEQ ID NO:1 of SEQUENCE LISTING. The amino acid sequence deduced from the above nucleotide sequence is shown under SEQ ID NO:2. Here, a sequence characteristic of a gene coding for decaprenyl diphosphate synthase is observed.

[0020] The DNA of the invention may be any of the DNA having the nucleotide sequence shown under SEQ ID NO:1, the DNA having a nucleotide sequence derived from the sequence of SEQ ID NO:1 by deletion, addition, insertion and/or substitution of one or a plurality of nucleotides and coding for a protein having decaprenyl diphosphate synthase activity, and the DNA which hybridizes with the DNA having the nucleotide sequence of SEQ ID NO:1 under stringent conditions and codes for a protein having decaprenyl diphosphate synthase activity.

[0021] The "nucleotide sequence derived by deletion, addition, insertion and/or substitution of one or a plurality of nucleotides" means any nucleotide sequence derived by deletion, addition, insertion and/or substitution of a number of nucleotides of the order which can be deleted, added, inserted and/or substituted by the methods well known in the art, for example as described in, *inter alia*, Protein, Nucleic Acid, Enzyme, Supplemental Issue: Gene Amplification PCR Technology TAKKAJ 35 (17), 2951-3178 (1990) and Henry A. Erlich (ed.), PCR Technology (the translation edited by Ikunoshin Kato) (1990).

[0022] As used in this specification, the term "protein having decaprenyl diphosphate synthase activity" means a protein capable of synthesizing decaprenyl diphosphate in a yield of not less than 10%, preferably not less than 40%, more preferably not less than 60%, still more preferably not less than 80%, relative to the protein having the amino acid sequence shown under SEQ ID NO:2. Such yield measurements can be made by the technique which comprises reacting FDP (farnesyl diphosphate) and ¹⁴C-IPP (radiolabeled isopentenyl diphosphate) with the enzyme of interest, hydrolyzing the resulting ¹⁴C-DPP (decaprenyl diphosphate) with phosphatase, fractionating the hydrolysate by TLC, and assaying the amounts taken up in spots corresponding to the respective chain lengths (Okada et al., Eur. J. Bio-

chem., 255, 55 to 59).

[0023] The "DNA which hybridizes with the DNA having the nucleotide sequence of SEQ ID NO:1 under stringent conditions" means a DNA obtained by colony hybridization, plaque hybridization, Southern hybridization or the like hybridization techniques using the DNA having the nucleotide sequence of SEQ ID NO : 1 as the probe. Anyone skilled in the art may easily acquire the objective DNA by carrying out said hybridization according to the methods described in Molecular Cloning, 2nd Edition (Cold Spring Harbor Laboratory Press, 1989).

[0024] The protein of the present invention may have the amino acid sequence shown under SEQ ID NO:2 or an amino acid sequence derived from the amino acid sequence shown under SEQ ID NO:2 by deletion, addition, insertion and/or substitution of one or a plurality of amino acids and having decaprenyl diphosphate synthase activity.

[0025] "The amino acid sequence derived by deletion, addition, insertion and/or substitution of one or a plurality of amino acids" can be obtained by effecting such deletion, addition, insertion and/or substitution by the technology well known in the art, such as region-specific mutagenesis techniques. Specific procedures are described in Nucleic Acid Res. 10, 6487 (1982), Methods in Enzymology, 100, 448 (1983) and other literature.

[0026] The protein of the present invention preferably has an amino acid sequence showing a homology of not less than 60%, preferably not less than 70%, more preferably not less than 80%, still more preferably not less than 90%, further still more preferably not less than 95%, to the amino acid sequence shown under SEQ ID NO:2.

[0027] The "homology" is calculated by aligning two nucleotide sequences to be compared in the optimum format, counting the matched base positions (A, T, C, G, U or I) between the two sequences, dividing the count by the total number of bases compared, and multiplying the product by 100. Specifically, this calculation can be made using an analytical software such as Hitachi Soft Engineering's DNASIS, Software Development's GENETYX, or Finland CSC's Clustal X, for instance.

[0028] While the gene coding for decaprenyl diphosphate synthase must be ligated downstream of a suitable promoter for expression, an expression vector can be constructed, for example by excising a DNA fragment containing the gene with a restriction enzyme or amplifying the enzyme-encoding gene selectively by PCR, followed by cloning it into a vector having a promoter. In the present invention, the expression vector into which the DNA coding for the protein having decaprenyl diphosphate synthase activity may be inserted is not particularly restricted but may for example be one constructed by ligating a suitable promoter to a plasmid derived from *E. coli*. The plasmid of *E. coli* origin includes pBR322, pBR325, pUC19 and pUC119, while the promoter includes T7 promoter, trp promoter, tac promoter, lac promoter and λ PL promoter. Further, as the expression vector of this invention, pGEX-2T, pGEX-3T, pGEX-3X (all from Pharmacia), pBluescript, pUC19 (from Toyobo), pMALC2, pET-3T and pUCNT (described in WO 94/03613), etc. can also be mentioned. Among these, pUCNT can be used with advantage. To mention a specific example, the vector pNTSa1 for the expression of a decaprenyl diphosphate synthase gene can be constructed by inserting the gene having the DNA sequence shown under SEQ ID NO:1 into the expression vector pUCNT.

[0029] Then, this enzyme gene expression vector is introduced into a suitable microorganism, whereby the microorganism is rendered capable of producing coenzyme Q_{10} . The host microorganism is not particularly restricted but *Escherichia coli* can be used with advantage. The *Escherichia coli* is not particularly restricted but includes such strains as XL1-Blue, BL-21, JM109, NM522, DH5 α , HB101 and DH5, among others. Among these, *E. coli* DH5 α can be used with particular advantage. For example, when the expression vector pNTSa1 containing the decaprenyl diphosphate synthase gene is introduced into this *E. coli* strain, the coenzyme Q_{10} , which the intact *E. coli* inherently does not produce, can be produced in a large amount. This *E. coli* DH5 α (pNTSa1) has been deposited with National Institute of Bioscience and Human-Technology (Higashi 1-1-3, Tsukuba-shi, Ibaraki, Japan) under the accession number of FERM BP-6844.

[0030] Furthermore, *Escherichia coli* KO229 (Journal of Bacteriology, 179, 3058-3060 (1997), the octaprenyl diphosphate synthase gene-knockout *E. coli* strain constructed by Kawamukai et al. as the host microorganism, is incapable of producing coenzyme Q_8 and can be utilized as the host for higher production of coenzyme Q_{10} .

[0031] The gene can be used not only singly but may be introduced together with another biosynthesis-related gene into a microorganism to thereby obtain still more satisfactory results.

[0032] Coenzyme Q_{10} can be produced by culturing the transformant obtained according to the invention and harvesting the product coenzyme Q_{10} in a *per se* known manner. When the host microorganism is a strain of *Escherichia coli*, either LB broth or M9 broth containing glucose and casamino acids can be used as the culture broth. In order that the promoter may be allowed to function with efficiency, the broth may be supplemented with a certain chemical such as isopropyl-thiogalactoside or indolyl-3-acrylic acid. Culture can be carried out at 37 °C for 17 to 24 hours, for instance, optionally under aeration or agitation. In the practice of the invention, the product coenzyme Q_{10} may be used after purification or as it is in the crude form, depending on the intended use. Isolation of coenzyme Q_{10} from the culture can be made-by using known separation and purification procedures in a suitable combination. As such known separation and purification procedures, there can be mentioned techniques utilizing solubilities, such as salting-out and solvent precipitation; techniques chiefly utilizing differences in molecular weight, such as dialysis, ultrafiltration, gel filtration and (SDS-)polyacrylamide gel electrophoresis; techniques utilizing differences in charge, such as ion ex-

change chromatography; techniques utilizing specific affinity, such as affinity chromatography; techniques utilizing differences in hydrophobicity, such as reversed-phase high performance liquid chromatography; and techniques utilizing differences in isoelectric point, such as isoelectric focusing, among others.

[0033] The use for the coenzyme Q₁₀ obtained according to the invention is not particularly restricted but the enzyme can be applied to pharmaceuticals with advantage.

BRIEF DESCRIPTION OF THE DRAWINGS

[0034] Fig. 1 is a restriction map of the pNTSa1 vector containing the decaprenyl diphosphate synthase gene.

[0035] Fig. 2 is an HPLC chart of the coenzyme Q₁₀ produced by the recombinant *Escherichia coli* DH5α as transformed with the decaprenyl diphosphate synthase gene.

[0036] Fig. 3 is an HPLC chart of the coenzyme Q₁₀ produced by the recombinant *Escherichia coli* KO229 as transformed with the decaprenyl diphosphate synthase gene.

BEST MODE FOR CARRYING OUT THE INVENTION

(Example 1)

[0037] The chromosome DNA of *Saitoella complicata* IFO 10748 was prepared by the method of C. S. Hoffman et al. (Gene, 57 (1987), 267-292). Based on the homology to the known long-chain prenyl diphosphate synthase genes, PCR primers, i.e. DPS-1 (5'-AAGGATCCTNYTNCAYGAYGT-3') and DPS-1 1AS (5'-ARYTGNADRAAYTCNCC-3'), were designed. In the above sequences, R stands for A or G; Y for C or T; and N for G, A, T or C. Using these primers, PCR was carried out under the conditions of heat treatment at 94 °C, 3 min. followed by 40 cycles of 94 °C, 1 min. → 43 °C, 2 min. → 72 °C, 2 min., and the PCR product was analyzed by 1.2% agarose gel electrophoresis.

[0038] The ca. 220 bp fragment thus obtained was excised from the gel and purified using a DNA extraction kit (Sephaglas™ BrandPrep Kit, Amersham Pharmacia Biotech). Then, using a PCR product direct cloning kit (pT7BlueT-Vector Kit, NOVAGEN), the DNA was cloned into the *E. coli* expression vector to give pT7-SaDPS. Then, using a DNA sequencer (Model 377, Perkin-Elmer) and a DNA sequencing kit (Perkin-Elmer; ABI PRISM™ BigDye™ Terminator Cycle Sequence Ready Reaction Kit with AmptiTaq™ DNA Polymerase, FS), DNA sequencing was carried out according to the kit manufacturer's protocol. As a result, there was obtained a sequence corresponding to the nucleotides 717 through 924 of SEQ ID NO:1 under SEQUENCE LISTING. The translation sequence thus obtained contained "GDFLLGRA" which is a characteristic region of polyprenyl diphosphate synthases and, therefore, was considered to be part of the decaprenyl diphosphate synthase gene.

(Example 2)

[0039] Using 0.03 µg of a pT7-SaDPS vector containing a 220 bp DNA fragment which was considered to be the decaprenyl diphosphate synthase gene of *Saitoella complicata* IFO 10748, PCR using primers Sa-1S (which has the sequence of 5'-GAGACCAGACGAAACGCACCA-3') and Sa-2AS (which has the sequence of 5'-TGGTGCCTT-TCGTCTGGTCTC-3') was carried out [94 °C, 3 min. → (94 °C, 30 sec. → 55 °C, 30 sec. → 72 °C, 1 min.) × 25 cycles → 72 °C, 5 min. → 4 °C]. The PCR product was subjected to gel electrophoresis using 1.2% agarose (Takara) and a ca. 145 bp fragment was excised from the gel and purified using a DNA extraction kit (Sephaglas™ BrandPrep Kit; Amersham Pharmacia Biotech). Using about 100 ng of this DNA fragment, chemiluminescence labeling was performed using ECL Direct Nucleic Acid Labeling System (Amersham Pharmacia Biotech).

(Example 3)

[0040] The chromosome DNA of *Saitoella complicata* IFO 10748 was digested with the restriction enzyme EcoRI and electrophoresed through 0.8% agarose gel. This gel was denatured with alkali (0.5 M NaOH, 1.5 M NaCl) and neutralized (0.5 M Tris-HCl (pH 7.5), 1.5 M NaCl), after which HYBOND N + filter (Amersham) was placed on the gel and the Southern transfer was carried out using 20 × SSC overnight. This filter was dried and heated at 80 °C for 2 hours and using ECL Direct Nucleic Acid Labeling/Detection System (Amersham Pharmacia Biotech), Southern hybridization and detection were carried out. Thus, using Gold Hybridization Solution (Amersham Pharmacia Biotech), prehybridization was performed at 42 °C for 1 hour.

[0041] The chemiluminescence-labeled probe was heated at 95 °C for 5 minutes, quenched on ice, and added to the prehybridization solution used for filter prehybridized and the hybridization was carried out at 42 °C for 22 hours. This filter was washed with 0.5 × SSC solution containing 6 M urea and 0.4% SDS at 42 °C twice for 20 minutes each and, then, washed with 2 × SSC solution at room temperature twice for 5 minutes each. This filter was immersed in

Enhanced Chemiluminescence Reagent (product of Amersham Pharmacia Biotech) and, then, exposed in intimate contact with X-ray film to detect a black exposure band.

[0042] As a result, the probe was found to have firmly hybridized with a ca. 10 kbp EcoRI restriction fragment.

(Example 4)

[0043] The chromosome DNA of *Saitoella complicata* IFO 10748 was digested with the restriction enzyme EcoRI and electrophoresed through 0.8% agarose and a ca. 10 kbp fragment of the DNA was excised from the gel and purified to prepare a DNA fragment for cloning. Using λ -DASHII Phage Kit (product of Stratagene), the above DNA fragment was inserted into the EcoRI site of its phage and the packaging was made using In Vitro Packaging Kit (Amersham). *Escherichia coli* XL1-Blue MRA (P2) was infected and layered on NZY plate medium (5g/L NaCl, 2 g/L $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 5 g/L yeast extract, 10 g/L NZ amine, 18 g/L agar (pH 7.5)) together with NZY soft agar (the agar only of NZY plate medium, 8 g/L) for use as a plaque. This was transferred to HYBOND N + filter (product of Amersham), denatured with alkali (0.5 M NaOH, 1.5 M NaCl), neutralized (0.5 M Tris-HCl (pH 7.5), 1.5 M NaCl), dried, and heated at 80 °C for 2 hours.

[0044] Using 9 filters heated as above, the prehybridization and the hybridization using the chemiluminescence-labeled probe were carried out as in Example 3 and the filters were rinsed. Each filter was then dried and exposed in intimate contact with X-ray film and the phage plaque corresponding to the black exposure spot was separated. The phage of the separated plaque was used to infect *E. coli* in the same manner as above and transferred to the filter and the hybridization was carried out again for confirmation. As a result, 6 phage clones could be selected.

[0045] Using a suspension of the phage, PCR was carried out using said Sa-1S and Sa-2AS primers, and as a result, a 145-bp DNA fragment could be detected in 6 clones. Therefore, the phage DNA was prepared from the recombinant λ -DASHII phage particles according to Laboratory Manual for Genetic Engineering (Masami Muramatsu, Maruzen, 1990). For subcloning, the phage DNA thus prepared was digested with the restriction enzymes Sall and SacI and electrophoresed through 0.8% agarose gel. This gel was denatured with alkali (0.5 M NaOH, 1.5 M NaCl) and neutralized (0.5 M Tris-HCl (pH 7.5), 1.5 M NaCl). Then, a HYBOND N + filter (Amersham) was placed on the gel and subjected to Southern transfer using 20 x SSC overnight. This filter was dried and heated at 80°C for 2 hours, after which the Southern hybridization and detection were carried out using ECL Direct Nucleic Acid Labeling/Detection System (Amersham Pharmacia Biotech). Thus, using Gold Hybridization Solution (Amersham Pharmacia Biotech), the prehybridization was carried out at 42 °C for 1 hour.

[0046] The chemiluminescence-labeled probe was heated at 95 °C for 5 minutes, quenched on ice, and added to the prehybridization solution used for filter prehybridized and the hybridization was carried out at 42 °C for 22 hours. This filter was washed with 0.5 x SSC containing 6 M urea and 0.4% SDS at 42 °C twice for 20 minutes each and, then, with 2 x SSC at room temperature twice for 5 minutes each. The filter was immersed in Enhanced Chemiluminescence Reagent (Amersham Pharmacia Biotech) and exposed in intimate contact with X-ray film to detect a black exposure band.

[0047] As a result, the probe was found to have intimately hybridized with a ca. 4.5 kb fragment as obtained by digestion with the restriction enzyme Sall and a ca. 3.5 kb fragment as obtained by digestion with SacI.

[0048] The phage DNA was digested with the restriction enzymes Sall and SacI and electrophoresed through 0.8% agarose gel. The restriction fragment corresponding to the position and size of the black exposure band was excised from the gel and purified using a DNA extraction kit (Sephaglas™ Brand Prep Kit; Amersham Pharmacia Biotech). Then, using a DNA sequencer (Model 377, Perkin-Elmer Corp.) and a DNA sequence kit (Perkin-Elmer Corp., ABI PRISM™ BigDye™ Terminator Cycle Sequence Ready Reaction Kit with AmpTaq™ DNA polymerase, FS), the sequencing was carried out in accordance with the manufacturer's protocol.

[0049] As a result, it was found that the Sall site and SacI site are located at positions 1124 and 1241, respectively, of SEQ ID NO:1 under SEQUENCE LISTING and that no fragment contained the C-terminal. So, for Sall, which is the upstream one of the two restriction enzymes, in the decaprenyl diphosphate synthase gene, the remaining fragments were examined. As a result, a 3 kbp fragment was found to contain a sequence including a terminal region of the SacI fragment up to the termination codon. By analyzing these 3 restriction fragments, the full-length sequence of the decaprenyl diphosphate synthase gene could be elucidated. Of the three DNA fragments, the ca. 1.6 kbp fragment was sequenced. The result is shown as SEQ ID NO:1 under SEQUENCE LISTING. Moreover, the amino acid sequence deduced from the above DNA sequence is shown as SEQ ID NO:2.

[0050] Comparison of the DNA sequence thus obtained with that of the decaprenyl diphosphate synthase gene of *Saccharomyces cerevisiae* as described in Journal of Biological Chemistry, 265, 13157-13164 (1990) revealed about 48% homology on the amino acid level as analyzed using Hitachi Soft Engineering's DNASIS software. Comparison with the decaprenyl diphosphate synthase derived from *Schizosaccharomyces pombe* as described in Japanese Kokai Publication Hei-9-173076 by means of DNASIS revealed 49% homology on the amino acid level.

(Example 5)

[0051] In order to selectively excise the gene region coding for decaprenyl diphosphate synthase from the prepared phage DNA, PCR was carried out using synthetic DNA primers Sa-N1 (which has the sequence of 5'-AACATATGGCCT-CACCAGCACTGCGG-3') and Sa-C (which has the sequence of 5'-AAGAATTCCTATCTTGACCTAGTCAACAC-3') in otherwise the same manner as in Example 3. After digestion with the restriction enzymes NdeI and EcoRI, the fragment was inserted into the expression vector pUCNT (disclosed in WO 94/03613) to construct the decaprenyl diphosphate synthase gene expression vector pNTSa1. The restriction map of the expression vector pNTSa1 thus obtained is shown in Fig. 1. It is to be noted that DPS represents the coding region of the decaprenyl diphosphate synthase gene.

(Example 6)

[0052] The decaprenyl diphosphate synthase gene expression vector pNTSa1 constructed as above was introduced into *Escherichia coli* DH5 α . The microorganism was shake-cultured in 10 mL of LB broth at 37°C overnight and the cells were harvested by centrifugation (3000 rpm, 20 min.).

[0053] The cells were suspended in 1 mL of 3% aqueous solution of sulfuric acid and heat-treated at 120°C for 30 minutes. Then, 2 mL of 14% aqueous solution of sodium hydroxide was added and the mixture was further heat-treated at 120 °C for 15 minutes. The lysate obtained was extracted with 3 mL of hexane-isopropyl alcohol (10:2), and after centrifugation, 1.5 mL of the organic layer was separated and evaporated to dryness under reduced pressure. The residue was dissolved in 200 μ L of ethanol and a 20 μ L portion of the solution was subjected to HPLC analysis (LC-10A, Shimadzu Corporation). Fractionation was carried out using a reversed-phase column (YMC-pack ODS-A, 250 \times 4.6 mm, S-5 μ m, 120 A) and, as the mobile phase, ethanol-methanol (2:1) and the coenzyme Q₁₀ produced was detected from the absorbance at the wavelength of 275 nm. The result is shown in Fig. 2. As can be seen from Fig. 2, it was found that when the decaprenyl diphosphate synthase gene is introduced into a host and allowed to be expressed, the resulting recombinant *Escherichia coli* produces coenzyme Q₁₀ which *E. coli* in general inherently does not produce.

[0054] The recombinant *E. coli* DH5 α (pNTSa1) obtained as above has been deposited with National Institute of Bioscience and Human-Technology (Higashi 1-1-3, Tsukuba-shi, Ibaraki, Japan) as of Heisei 11, August 17 (accession number FERM BP-6844).

(Example 7)

[0055] The octaprenyl diphosphate synthase gene-knockout *Escherichia coli* KO229 constructed by Kawamukai et al. is known to retain the gene supported on the spectinomycin-resistant plasmid (pKA3) and die on dropout of said plasmid (Journal of Bacteriology, 179, 3058-3060 (1997)). The pNTSa1 was introduced into the above knockout strain, culturing the microorganism in 10 mL of ampicillin-containing LB broth by shake culture at 37 °C overnight, subculturing 1% of the culture in 10 mL of fresh ampicillin-containing LB broth and culturing the microorganism further by shake-culture at 37 °C overnight, and after 9 cycles of the above cultural procedure, selecting the strain growing on ampicillin-containing LB plate medium but not growing on spectinomycin-containing LB plate medium.

(Example 8)

[0056] The pNTSa1-transfected strain constructed in Example 7 was shake-cultured in 10 mL of LB broth at 37 °C overnight and the cells were harvested by centrifugation (3000 rpm, 20 min.).

[0057] The cells were suspended in 1 mL of 3% aqueous solution of sulfuric acid and heat-treated at 120 °C for 30 min. Then, 2 mL of 14% aqueous solution of sodium hydroxide was added and the mixture was further heat-treated at 120 °C for 15 minutes. The lysate cells were extracted with 3 mL of hexane-isopropyl alcohol (10:2) and centrifuged to recover 1.5 mL of the organic layer and the solvent was evaporated to dryness under reduced pressure. The residue was dissolved in 200 μ L of ethanol and a 20 μ L portion of the solution was subjected to HPLC analysis (LC-10A, Shimadzu Corporation). Fractionation was carried out by using a reversed-phase column (YMC-pack ODS-A, 250 \times 4.6 mm, S-5 μ m, 120 A) and, as the mobile phase, ethanol-methanol (2:1) and the product coenzyme Q₁₀ was detected from the absorbance at the wavelength of 275 nm. The result is shown in Fig. 3. It is clear from Fig. 3 that as the decaprenyl diphosphate synthase gene was introduced and allowed to be expressed, the host *Escherichia coli* was enabled to produce coenzyme Q₁₀ which it inherently does not produce and enabled to be transformed so as to yield coenzyme Q₁₀ in an increased amount more than that of the coenzyme Q₈ producer *E. coli* strain.

INDUSTRIAL APPLICABILITY

[0058] The gene coding for the key enzyme associated with the biosynthesis of coenzyme Q₁₀, namely decaprenyl diphosphate synthase, was isolated from fungi of the genus Saitoella and its nucleotide sequence was elucidated. Furthermore, the gene was successfully introduced and expressed in Escherichia coli. By utilizing the technology of the invention, coenzyme Q₁₀ in use as a pharmaceutical can be produced with improved efficiency.

10

15

20

25

30

35

40

45

50

55

Sequence listing

5 <110> Kaneka Corporation

<120> Process for producing coenzyme Q₁₀

10 <130> T549/QX-GT2

<150> JP P1999-237561

15 <151> 1999-08-24

<160> 2

20 <210> 1

<211> 1653

<212> DNA

25 <213> Saioella complicata

<400> 1

30 ttttgtgggg tcgaaaagtc ggcacgggtg caggttcggc ttgagaccag taaaggctcg 60

gagattgagt tcaggacaaa gctttgatcc gtgaggtcta catcttcagc aaatcatttc 120

35 aaatccatat acc atg gcc tca cca gca ctg cgg ata cga agc atc agc 169

Met Ala Ser Pro Ala Leu Arg Ile Arg Ser Ile Ser

1 5 10

40 tct cga tca atc gcc tct ctg cga tcg gtt acc cta aga aca gcc tcg 217

Ser Arg Ser Ile Ala Ser Leu Arg Ser Val Thr Leu Arg Thr Ala Ser

15 20 25

45 gca cct tca tta cga cta aga tgt acc ccg acg agc cgg cca tcg agt 265

Ala Pro Ser Leu Arg Leu Arg Cys Thr Pro Thr Ser Arg Pro Ser Ser

50 30 35 40

tca tgg gct gct gct gtg tct tcg gcg tcg aga ctg gtt gag cct gat 313

Ser Trp Ala Ala Ala Val Ser Ser Ala Ser Arg Leu Val Glu Pro Asp

55 45 50 55 60

55

	gag acc aga cga aac gca cca tcc gga aac cag gca ttc gga aac aag	793
	Glu Thr Arg Arg Asn Ala Pro Ser Gly Asn Gln Ala Phe Gly Asn Lys	
5	205 210 215 220	
10	atg gcg att ttg gct ggt gat ttc ttg ttg gga cgg gcg tct gtt gca	841
	Met Ala Ile Leu Ala Gly Asp Phe Leu Leu Gly Arg Ala Ser Val Ala	
	225 230 235	
15	ttg gcg agg ttg cgc aat ccg gag gtg att gag ctt ttg gct act gtt	889
	Leu Ala Arg Leu Arg Asn Pro Glu Val Ile Glu Leu Leu Ala Thr Val	
	240 245 250	
20	att gca aac ttg gtt gag gga gag ttc atg cag ttg aaa aat act gtt	937
	Ile Ala Asn Leu Val Glu Gly Glu Phe Met Gln Leu Lys Asn Thr Val	
	255 260 265	
25	gat gat gcg att gag gct acg gcg acg cag gaa acg ttc gat tac tat	985
	Asp Asp Ala Ile Glu Ala Thr Ala Thr Gln Glu Thr Phe Asp Tyr Tyr	
30	270 275 280	
	ttg cag aag act tac ttg aag act gcg tcc ttg att gcc aag tcg tgc	1033
	Leu Gln Lys Thr Tyr Leu Lys Thr Ala Ser Leu Ile Ala Lys Ser Cys	
35	285 290 295 300	
40	aga gca agt gcg ctt ctg ggt ggt gct acg cct gag gtt gct gat gct	1081
	Arg Ala Ser Ala Leu Leu Gly Gly Ala Thr Pro Glu Val Ala Asp Ala	
	305 310 315	
45	gct tat gct tac gga agg aac ctt ggt ttg gca ttc cag atc gtc gac	1129
	Ala Tyr Ala Tyr Gly Arg Asn Leu Gly Leu Ala Phe Gln Ile Val Asp	
	320 325 330	
50	gac atg ctc gac tac acc gtc tcc gct acc gac ctc ggt aag ccc gcc	1177
	Asp Met Leu Asp Tyr Thr Val Ser Ala Thr Asp Leu Gly Lys Pro Ala	
	335 340 345	

55

55

<210> 2

<211> 440

<212> PRT

<213> Saioella complicata

<400> 2

Met	Ala	Ser	Pro	Ala	Leu	Arg	Ile	Arg	Ser	Ile	Ser	Ser	Arg	Ser
1				5					10					15
Ile	Ala	Ser	Leu	Arg	Ser	Val	Thr	Leu	Arg	Thr	Ala	Ser	Ala	Pro
			20						25					30
Ser	Leu	Arg	Leu	Arg	Cys	Thr	Pro	Thr	Ser	Arg	Pro	Ser	Ser	Ser
			35						40					45
Trp	Ala	Ala	Ala	Val	Ser	Ser	Ala	Ser	Arg	Leu	Val	Glu	Pro	Asp
			50						55					60
Pro	Asn	Gln	Pro	Leu	Ile	Asn	Pro	Leu	Asn	Leu	Val	Gly	Pro	Glu
			65						70					75
Met	Ser	Asn	Leu	Thr	Ser	Asn	Ile	Arg	Ser	Leu	Leu	Gly	Ser	Gly
			80						85					90
His	Pro	Ser	Leu	Asp	Thr	Val	Ala	Lys	Tyr	Tyr	Val	Gln	Ser	Glu
			95						100					105
Gly	Lys	His	Ile	Arg	Pro	Leu	Met	Val	Leu	Leu	Met	Ala	Gln	Ala
			110						115					120
Thr	Glu	Val	Ala	Pro	Lys	Val	Gln	Gly	Trp	Glu	Lys	Val	Val	Glu
			125						130					135
Val	Pro	Val	Asn	Glu	Gly	Leu	Ala	Pro	Pro	Glu	Val	Leu	Asn	Asp
			140						145					150
Lys	Asn	Pro	Asp	Met	Met	Asn	Met	Arg	Ser	Gly	Pro	Leu	Thr	Lys
			155						160					165
Asp	Gly	Glu	Ile	Glu	Gly	Gln	Thr	Ser	Asn	Ile	Leu	Ala	Ser	Gln
			170						175					180
Arg	Arg	Leu	Ala	Glu	Ile	Thr	Glu	Met	Ile	His	Ala	Ala	Ser	Leu
			185						190					195
Leu	His	Asp	Asp	Val	Ile	Asp	Ala	Ser	Glu	Thr	Arg	Arg	Asn	Ala
			200						205					210
Pro	Ser	Gly	Asn	Gln	Ala	Phe	Gly	Asn	Lys	Met	Ala	Ile	Leu	Ala
			215						220					225
Gly	Asp	Phe	Leu	Leu	Gly	Arg	Ala	Ser	Val	Ala	Leu	Ala	Arg	Leu
			230						235					240

	Arg Asn Pro Glu Val Ile Glu Leu Leu Ala Thr Val Ile Ala Asn	
	245	250 255
5	Leu Val Glu Gly Glu Phe Met Gln Leu Lys Asn Thr Val Asp Asp	
	260	265 270
	Ala Ile Glu Ala Thr Ala Thr Gln Glu Thr Phe Asp Tyr Tyr Leu	
10	275	280 285
	Gln Lys Thr Tyr Leu Lys Thr Ala Ser Leu Ile Ala Lys Ser Cys	
	290	295 300
15	Arg Ala Ser Ala Leu Leu Gly Gly Ala Thr Pro Glu Val Ala Asp	
	305	310 315
	Ala Ala Tyr Ala Tyr Gly Arg Asn Leu Gly Leu Ala Phe Gln Ile	
	320	325 330
20	Val Asp Asp Met Leu Asp Tyr Thr Val Ser Ala Thr Asp Leu Gly	
	335	340 345
	Lys Pro Ala Gly Ala Asp Leu Gln Leu Gly Leu Ala Thr Ala Pro	
	350	355 360
25	Ala Leu Phe Ala Trp Lys His His Ala Glu Leu Gly Pro Met Ile	
	365	370 375
	Lys Arg Lys Phe Ser Asp Pro Gly Asp Val Glu Arg Ala Arg Glu	
30	380	385 390
	Leu Val Glu Lys Ser Asp Gly Leu Glu Lys Thr Arg Ala Leu Ala	
	395	400 405
35	Glu Glu Tyr Ala Gln Lys Ala Leu Asp Ala Ile Arg Thr Phe Pro	
	410	415 420
	Glu Ser Pro Ala Arg Lys Ala Leu Glu Gln Leu Thr Asp Lys Val	
	425	430 435
40	Leu Thr Arg Ser Arg	
	440	

45

Claims

1. A DNA of the following (a), (b) or (c):

50

- (a) a DNA having the nucleotide sequence shown under SEQ ID NO:1;
- (b) a DNA having a nucleotide sequence derived from the nucleotide sequence of SEQ ID NO:1 by deletion, addition, insertion and/or substitution of one or a plurality of nucleotides and coding for a protein having decaprenyl diphosphate synthase activity;
- (c) a DNA which hybridizes with the DNA having the nucleotide sequence of SEQ ID NO:1 under stringent conditions and codes for a protein having decaprenyl diphosphate synthase activity.

55

2. A protein of the following (d) or (e):

(d) a protein having the amino acid sequence shown under SEQ ID NO:2;
 (e) a protein having an amino acid sequence derived from the amino acid sequence of SEQ ID NO:2 by deletion, addition, insertion and/or substitution of one or a plurality of amino acids and having decaprenyl diphosphate synthase activity.

5

3. A DNA coding for the protein according to Claim 2.

4. An expression vector constructed by cloning the DNA according to Claim 1 or 3 in an expression vector.

10

5. The expression vector according to Claim 4
 wherein the expression vector is pUCNT.

6. The expression vector according to Claim 5
 wherein the expression vector is pNTSa1.

15

7. A transformant as obtainable by transforming a host microorganism with the DNA according to Claim 1 or 3.

8. A transformant as obtainable by transforming a host microorganism using the expression vector according to Claim 4, 5 or 6.

20

9. The transformant according to Claim 7 or 8 wherein the host microorganism is Escherichia coli.

10. The transformant according to Claim 9 wherein the Escherichia coli is Escherichia coli DH5 α .

25

11. The transformant according to Claim 10 which is E. coli DH5 α (pNTSa1) (FERM BP-6844).

12. A process for producing a coenzyme Q₁₀, which comprises culturing the transformant according to Claim 7, 8, 9, 10 or 11 in a culture broth and harvesting the coenzyme Q₁₀ produced and accumulated in the resulting culture.

30

35

40

45

50

55

Fig. 1






Fig. 2

E. Coli DH5 α



E. Coli DH5 α / pNTSa1



CoQ₁₀ Standard






Fig. 3



E. Coli K0229 / pKA3



E. Coli K0229 / pKA3 + pNTSa1



E. Coli K0229 / pNTSa1

CoQ₁₀ Standard

INTERNATIONAL SEARCH REPORT

International application No.

PCT/JP00/05659

A. CLASSIFICATION OF SUBJECT MATTER Int.Cl ⁷ C12N15/54, C12N9/12, C12N1/21, C12P7/66		
According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) Int.Cl ⁷ C12N15/54, C12N9/12, C12N1/21, C12P7/66		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) SwissProt/PIR/GeneSeq, WPI (DIALOG), BIOSIS (DIALOG)		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	JP, 9-173076, A (Alpha Shokuhin K.K.), 08 July, 1997 (08.07.97) (Family: none)	1-12
A	Suzuki. K., et al., "Analysis of the Decaprenyl Diphosphate Synthase (dps) Gene in Fission Yeast Suggests a Role of Ubiquinone as an Antioxidant", J. Biochem. (1997) Vol.121, pp.496-505	1-12
<input type="checkbox"/> Further documents are listed in the continuation of Box C. <input type="checkbox"/> See patent family annex.		
* Special categories of cited documents: "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed		"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "&" document member of the same patent family
Date of the actual completion of the international search 12 December, 2000 (12.12.00)		Date of mailing of the international search report 26 December, 2000 (26.12.00)
Name and mailing address of the ISA/ Japanese Patent Office		Authorized officer
Facsimile No.		Telephone No.

Form PCT/ISA/210 (second sheet) (July 1992)